

Some extensions of the class of k -convex bodies

V. Golubyatnikov*

Sobolev Institute of Mathematics, Novosibirsk, 630090, Russia

e-mail: glbntn@math.nsc.ru

V. Rovenski† *University of Haifa, Haifa, 31905, Israel*

e-mail: rovenski@math.haifa.ac.il

Abstract.¹ We study relations of some classes of k -convex, k -visible bodies in Euclidean spaces. We introduce and study circular projections in normed linear spaces and classes of bodies related with families of such maps, in particular, k -circular convex and k -circular visible ones. Investigation of these bodies more general than k -convex and k -visible ones allows us to generalize some classical results of geometric tomography and find their new applications.

Introduction

As usual, a convex body in n -dimensional Euclidean space $\mathbb{E}^n$ (over reals, complex or quaternion numbers) is a compact convex K set with non-empty interior, $\text{int}(K)$, whose closure coincides with the body, $\text{cl}(\text{int}(K)) = K$. The classical **problem** of geometric tomography concerns the reconstructing a body K of $\mathbb{E}^n$ from the knowledge of its orthogonal projections onto $(n - k)$ -planes of a preselected family $\mathcal{P}_k$.

Convex bodies are characterized by the property that they can be represented as intersections of families of (supporting) half-spaces. The axiomatic approach to the notion of convexity consists of the following. In an arbitrary set X , we choose a certain family $\mathcal{B}$ of subsets (called the convexity base). A set K is called $\mathcal{B}$ -convex if it is an intersection of some subfamily of sets from $\mathcal{B}$.

Reshetnyak [1] considered some generalization of convex surfaces, called δ -touched surfaces, in order to study behavior of geodesic lines on these surfaces. Following the same direction we have introduced in [2] the ε -convex bodies (more general than convex ones) and investigated the problem of determination of ε -convex bodies by their projection-type images. Instead of supporting half-spaces (as a family $\mathcal{B}$) we used there some different shapes such as complements to balls and cylinders and introduced circular projections onto punctured hyperplanes.

*Supported by President of Russia grant for leading scientific schools No. 8526.2006.1 and grant FP7-PEOPLE-2007-2-1-IEF, No. 219696 of Marie-Curie action.

¹*Keywords and phrases:* k -convex, k -visible body, supporting ball, circular projections, visual hull, geometric tomography

AMS Mathematics Subject Classification: 52A01, 52A30, 52A35

Some results for the convex bodies can be proved for k -convex and k -visible bodies (see [3]) and k -visual bodies (see [4], [5]) as well. The definition of a k -convex submanifold was introduced in [6] as a multi-dimensional generalization of a convex surface. In order to prove stability of determination of k -visible bodies by their images under projections, the first author [7] introduced α -rough bodies, where $\alpha \in (0, \pi]$; for $\alpha = \pi$ we get the class of convex bodies.

In the present work we extend the classes of k -convex and k -visible bodies in $\mathbb{E}^n$ and study the problems of their reconstruction from the knowledge of orthogonal projections onto $(n - k)$ -planes. We introduce our main tool, the families $\mathcal{P}_k$ of **circular projections** onto punctured $(n - k)$ -dimensional planes.

In order to extend our previous results [2], [3], [8], [9], following to the "soft-hard" ranking of geometrical categories described in [10], we study the classes of $\mathcal{C}$ -**convex** $\mathcal{K}\mathcal{P}_i^k, \mathcal{K}_i^{k,\varepsilon}$ and $\mathcal{V}\mathcal{P}_i^k, \mathcal{V}_i^{k,\varepsilon}$ ($i = 1, 2, \varepsilon > 0$) of $\mathcal{C}$ -**visible** bodies, more general than ε -convex or ε -visible classes of bodies introduced in [2]. In the definitions of these classes we consider

- the boundary points of the bodies ($i = 1$),
- the points disjoint from the bodies ($i = 2$).

We prove strong inclusions between these classes. Note that the classes $\mathcal{K}_2^k, \mathcal{K}\mathcal{P}_2^k$ and $\mathcal{K}_2^{k,\varepsilon}$ are closed under intersections of bodies, some results for the second type of bodies ($i = 2$) can be extended to the wider 1st one ($i = 1$).

It is interesting to find conditions **C** such that "If the bodies $U, V \subset \mathbb{E}^n$ satisfy **C** and their circular projections of a preselected family $\mathcal{P}_k$ coincide then $U = V$ ".

Much more hard and general **question** is the following: What kind of condition **CT** and what kind of transformations **PT**, **ST** should be taken in order to get a proposition: "If $U, V \subset \mathbb{E}^n$ satisfy **CT** and their circular projections of a family $\mathcal{P}_k$ are **PT**-equivalent, then the bodies U, V are **ST**-equivalent in the ambient space"?

For the case of orthogonal projections some answers to the question above were obtained in [3], [8]. The study of circular projections and circular $\mathcal{C}$ -visible bodies allows us to extend some classical results of the theory of convex bodies, see for example [3], [11], and to find their new applications in geometric tomography.

1 k -Convex bodies and visual hulls of sets

In what follows we as usual consider compact bodies in $\mathbb{E}^n$ over reals. Let $P^k \subset \mathbb{E}^n$ be a k -dimensional plane (i.e., an affine k -subspace). Denote by

$$B(P^k, r) = \{x \in \mathbb{E}^n : \text{dist}(x, P^k) \leq r\} \quad (1)$$

a solid cylinder of radius $r > 0$ with the axis P^k , that is the metric product of $\mathbb{E}^k$ and a ball $B^{n-k}(O, r) \subset \mathbb{E}^{n-k}$. For $k = 0$, (1) gives us a ball $B^n(C, r)$.

Definition 1 Let L be a body (usually we consider balls or cylinders) or a k -dimensional plane in $\mathbb{E}^n$. We call L **supporting** a body K if L intersects ∂K and L is disjoint from $\text{int } K$.

A half-plane $P_{\geq}^{k+1}$ can be represented as the union $P_{>}^{k+1} \cup P^k$ of an open half-plane $P_{>}^{k+1}$ and a boundary k -plane. A half-plane $P_{\geq}^{k+1}$ will be called **supporting** a body K if $P^k \cap \partial K \neq \emptyset$ and $P_{>}^{k+1} \cap B(P^k, r)$ is disjoint from $\text{int } K$ for some $r > 0$ depended only on K . In this case, obviously, P^k is supporting K . In what follows for simplicity we assume $r = \infty$.

Definition 2 A body $K \subset \mathbb{E}^n$ ($n > 2$) is called **k -convex of a class $\mathcal{K}_i^k$** if

$\mathcal{K}_1^k$: any point $x \in \partial K$ belongs to the boundary of a half-plane $P_{\geq}^{k+1}$ supporting K .

$\mathcal{K}_2^k$: any point $x \notin K$ belongs to the boundary of a half-plane $P_{\geq}^{k+1}$ disjoint from K .

A body $K \subset \mathbb{E}^n$ is called **k -visible of a class $\mathcal{V}_i^k$** if

$\mathcal{V}_1^k$: any plane Q^{k-m} supporting K belongs to the boundary of a half-plane $P_{\geq}^{k+1}$ supporting K .

$\mathcal{V}_2^k$: any plane Q^{k-m} , which is disjoint from K , belongs to the boundary of a half-plane $P_{\geq}^{k+1}$ disjoint from K .

Here $0 < m \leq k < n$, and the 0-dimensional plane is a point.

A connected boundary component of a k -convex (or k -visible) body K is called a **k -convex (or k -visible) hypersurface** of a certain class listed above.

Remark 1 (a) A Π -shaped non-convex body in $\mathbb{E}^3$ belongs to $\mathcal{K}_1^k$.

(b) A connected body of a class $\mathcal{K}_1^{n-1}$ is a convex one.

Proposition 1 The conditions $\mathcal{K}_i^k$ ($i = 1, 2$) are equivalent to the next ones:

($i = 1$) for any $x \in \partial K$, $K \in \mathcal{K}_1^k$, there is orthogonal projection $f : \mathbb{E}^n \rightarrow P^{n-k}$ such that $f(x)$ is a vertex of a ray supporting $f(K)$.

($i = 2$) for any $x \notin K$, $K \in \mathcal{K}_2^k$, there is orthogonal projection $f : \mathbb{E}^n \rightarrow P^{n-k}$ such that $f(x)$ is a vertex of a ray disjoint from $f(K)$.

The condition $\mathcal{V}_i^k$ ($i = 1, 2$) is equivalent to the following one:

($i = 1$) for any plane Q^{k-m} supporting K , $K \in \mathcal{V}_1^k$, there is orthogonal projection $f : \mathbb{E}^n \rightarrow P^{n-k}$ such that $f(Q^{k-m})$ is a vertex of a ray supporting $f(K)$.

($i = 2$) for any plane Q^{k-m} disjoint from K , $K \in \mathcal{V}_2^k$, there is orthogonal projection $f : \mathbb{E}^n \rightarrow P^{n-k}$ such that $f(Q^{k-m})$ is a vertex of a ray disjoint from $f(K)$.

The proof of Proposition 1 follows from the definitions.

Remark 2 The classes of **k -convex** and **k -visible** bodies can be defined relative to any non-empty family $\mathcal{P}_k$ of orthogonal projections onto $(n - k)$ -planes.

Class $\mathcal{V}_2^k$ for $k = 1$ is closed under the intersection of bodies in $\mathbb{E}^n$, for $k > 1$ this claim fails, [3]. Note that $\mathcal{V}_i^1 = \mathcal{K}_i^1$ ($i = 1, 2$). Remark that the properties ($i = 1$) fail if one will replace the definitions of $\mathcal{K}_1^k$ and $\mathcal{V}_1^k$ by weaker ones:

$\mathcal{K}_{1-}^k$: any point $x \in \partial K$ belongs to a k -plane P^k supporting K .

$\mathcal{V}_{1-}^k$: any plane Q^{k-m} supporting K belongs to a plane P^k supporting K as well.

Similarly one may formulate weaker conditions $\mathcal{K}_{2-}^k$ and $\mathcal{V}_{2-}^k$. Some of results of this section can be proved for the above classes of bodies.

Example 1 (a) The union of black squares of a chess-board and its n -dimensional analogues provide examples of bodies (not homeomorphic to a ball) in $\mathbb{E}^n$ of the classes $\mathcal{K}_1^{n-1} \setminus \mathcal{K}_1^{n-1}$ and $\mathcal{V}_1^{n-1} \setminus \mathcal{V}_1^{n-1}$. Example with homeomorphic to a ball bodies is given in what follows.

(b) Consider a homeomorphic to a ball body K in $\mathbb{E}^3$ between two helicoids, surfaces M_1 and M_0 , where $M_h : [u \cos v, u \sin v, v + h]$ and $0 \leq v \leq 2\pi, 0 \leq u \leq 1$. The projection of K onto xy -plane is a unit disc. The property $(\mathcal{K}_1^k)$ for $k = 1$ is satisfied, but the xy -projection of the z -axis (supporting K) is the inner point (the origin) of the disc. Hence, $K \notin \mathcal{K}_1^k$.

The following result was proved in [3] for a smaller class $\mathcal{V}_2^k$.

Lemma 1 (a) A connected body $K \in \mathcal{V}_1^{n-1}$ in $\mathbb{E}^n$ is convex. (b) The projection of a body $K \in \mathcal{V}_1^k$ in $\mathbb{E}^n$ onto any hyperplane belongs to a class $\mathcal{V}_1^{k-1}$. The corresponding visibility will be for its projections onto planes of smaller dimension, in particular, the projection of a body $K \in \mathcal{V}_1^{n-2}$ in $\mathbb{E}^n$ onto any 3-plane belongs to a class $\mathcal{V}_1^1$.

Proof. Part (a) of lemma follows from the fact that the body K lies on one side of its supporting hyperplane.

(b) Let $K(\omega)$ be the projection of K onto a hyperplane $P^{n-1}(\omega)$, orthogonal to unit vector $\omega \in \mathbb{E}^n$ and let $y \in \partial K(\omega)$. All inverse images of this point under projecting lie on the boundary of K . Let $z \in \partial K$ be one of inverse images of the point y . The line $l(yz)$ is supporting K and, by definition of the class $\mathcal{V}_1^k$, belongs to $(k+1)$ -dimensional half-plane H^{k+1} , supporting K . The image of H^{k+1} under projection onto $P^{n-1}(\omega)$ is a k -dimensional half-plane, supporting $K(\omega)$. $\square$

Theorem 1 $\mathcal{K}_2^k \subset \mathcal{K}_1^k$, $\mathcal{V}_2^k \subset \mathcal{V}_1^k$ and $\mathcal{V}_i^k \subset \mathcal{K}_i^k$ ($i = 1, 2$).

Proof. $\mathcal{K}_2^k \subset \mathcal{K}_1^k$. Given $K \in \mathcal{K}_2^k$ and $\bar{x} \in \partial K$, consider a sequence $x_i \notin K$ such that $x_i \rightarrow \bar{x}$ for $i \rightarrow \infty$. Due to $(\mathcal{K}_2^k)$, for each i there is a k -plane P_i^k containing x_i which bounds a half-plane $P_{\geq, i}^{k+1}$ disjoint from K . Let p_i be the normal to P_i^k at x_i directed inside of $P_{\geq, i}^{k+1}$. Taking a subsequence of integers, one may assume that there are a unit vector $\bar{p} = \lim_{i \rightarrow \infty} p_i$ and a k -plane $\bar{P}^k = \lim_{i \rightarrow \infty} P_i^k$ containing $\bar{x}$. Then $\bar{P}^k$ bounds a half-plane $\bar{P}_{>}^{k+1}$ and $\bar{p} \perp \bar{P}^k$ is a unit normal to $\bar{P}^k$ at $\bar{x}$ directed inside of $\bar{P}_{\geq}^{k+1}$. We claim that $\bar{P}_{\geq}^{k+1}$ is supporting K . To show this assume an opposite, that is $P_{>}^{k+1}$ contains a point $y \in \text{int } K$. Using standard isometry $\phi_i : \bar{P}_{\geq}^{k+1} \rightarrow P_{\geq, i}^{k+1}$, $\phi_i(\bar{x}) = x_i$, we find a sequence of points $y_i = \phi_i(y) \notin K$ converging to y . From this it follows that $y \notin \text{int } K$, that is a contradiction. Other inclusions can be proved analogously. Strong inclusions follow from examples in the sequel. $\square$

Example 2 (a) $K \in \mathcal{K}_1^k \setminus \mathcal{K}_2^k$ for $k = 1, n = 3$. Consider an envelope D (homeomorphic to a disc) with four unit square faces, and two sections along p and q , see Fig. 1. Let us bend $D \times [-c, c]$ (for small c) about r by an angle α




Figure 1: A body $K \in \mathcal{K}_1^k \setminus \mathcal{K}_2^k$ ($k = 1$), its "envelope" and model of 13 cubes.

bit smaller than 90° , and by 90° about the lines p, q . We obtain a body $K \subset \mathbb{E}^3$ that partially surrounds a cube (from four faces). Obviously, $K \in \mathcal{K}_1^1$, but the property $\mathcal{K}_2^1$ is not satisfied for the point $C(0, -1/4, 1/4) \notin K$ close to the front face (r). A cylinder $K' = K \times [-a, a] \subset \mathbb{E}^4$ is the body $K' \in \mathcal{V}_1^k \setminus \mathcal{V}_2^k$ for $k = 2, n = 4$.

(b) Example of a body $K \in \mathcal{K}_2^k \setminus \mathcal{V}_2^k$ for $k = 2, n = 4$, (i.e., a 2-convex body in $\mathbb{E}^4$ that is not 2-visible) is given in [3].

The following simple result was obtained by the first author on the initial stage of the study of the question that was formulated in Introduction.

Lemma 2 ([3]) *Let $V_1, V_2 \in \mathcal{K}_2^k$ be compact bodies in $\mathbb{E}^n, n > 2, n - k > 1$ (real, complex or quaternion) and G be a group of all either translations, or homotheties or is trivial. If the projections of V_1, V_2 onto any $(n - k)$ -plane P coincide relative to G , then V_1, V_2 also coincide relative to G .*

This claim is wrong in the real case for $n - k = 1$ and non-trivial G , since two different bodies of constant width have isometric projections onto any line.

Remark 3 Let $K \in \mathcal{K}_1^k$ ($k > n/2$) be a body in $\mathbb{E}^n$, and let ∂K be a C^2 -regular hypersurface with a unit normal ξ directed inside. By conditions, any point $x \in \partial K$ belongs to a plane P_x^k disjoint from $\text{int } K$. From this we conclude that $P_x^k \subset (TM)_x$, and the 2nd fundamental form of ∂K is non-negative on P_x^k . Hence at least k eigenvalues of the (symmetric) Weingarten operator A_ξ are non-negative. In this case, the homology groups (with integer coefficients) $H_{n-1-k+s}(\partial K) = 0$ for $0 < s < 2k - n - 1$, see Lemma 16 in [12] and also Lemma 6(b) in [9].

Definition 3 Given set W in $\mathbb{E}^n$ and a non-empty family $\mathcal{P}_k$ (of orthogonal projections onto $(n - k)$ -planes), a **visual hull** $\mathcal{P}_k\langle W \rangle$ is the largest set V such that $f(V) \subseteq f(W)$ for all $f \in \mathcal{P}_k$. A body $K \subset \mathbb{E}^n$ is **visual** relative to $\mathcal{P}_k$ if $\mathcal{P}_k\langle K \rangle = K$ [4].

If f is a projection onto subspace $P \subset \mathbb{E}^n$, $\mathcal{R}(f)$ denotes its range, i.e., P . Denote by $\bar{\mathcal{P}}_k$ a family of orthogonal projections onto all $(n - k)$ -planes in $\mathbb{E}^n$.

Proposition 2 ([4]) $\mathcal{P}_k\langle K \rangle = K$ if and only if for every $x \notin K$ there exists a $f \in \mathcal{P}_k$ such that $x + \mathcal{R}(f)^\perp$ does not intersect K .

The **visual hulls** of bodies in $\mathbb{E}^3$ for finite families $\mathcal{P}_k$ (reconstructions of 3-D shapes from multiple calibrated photographs) are studied in computational geometry and computer vision, see [5], [13], etc.

Proposition 3 $K \in \mathcal{K}_2^k$ if and only if $\bar{\mathcal{P}}_{n-k}\langle K \rangle = K$.

Proof. Suppose that there is $x \in \bar{\mathcal{P}}_{n-k}\langle K \rangle \setminus K$. By definition of $\mathcal{K}_2^k$, x belongs to some k -plane P^k which is disjoint from K , hence the projection of K onto orthogonal $(n-k)$ -plane $(P^k)^\perp$ does not contain the image of x . Thus $x \notin \bar{\mathcal{P}}_{n-k}\langle K \rangle$, a contradiction. $\square$

2 Circular projections and convexity

We will extend the classes of k -convex bodies in $\mathbb{E}^n$ in a way that the role of k -planes will play the k -dimensional spheres of radius $1/\varepsilon$ for a given $\varepsilon > 0$, and the circular projections will be used instead of orthogonal ones.

2.1 Circular projections

A $(n-k)$ -plane $P(C)$ containing a point $C \in \mathbb{E}^n$ is called the **punctured $(n-k)$ -plane**. Denote by $P^\perp(C)$ the punctured k -plane orthogonal to $P(C)$. Let $S^{n-1}(C, r)$ be a sphere of radius $r > 0$ centered in C , and $S^{n-k-1}(P(C), r) = S^{n-1}(C, r) \cap P(C)$ be a sphere of the same radius and center. Given $x \notin P^\perp(C)$, let $S^\perp(P(C), x)$ be a large k -sphere in a sphere $S^{n-1}(C, |Cx|)$ through x and orthogonal to a large sphere $S^{n-k-1}(P(C), |Cx|)$.

Definition 4 Let $P(C)$ be a punctured $(n-k)$ -plane. Define a **circular projection** $f_{P(C)} : \mathbb{E}^n \setminus P^\perp(C) \rightarrow P(C) \setminus C$ as follows. Given $x \notin P^\perp(C)$, let $f_{P(C)}(x)$ be the closest to x point of the sphere $S^{n-k-1}(P(C), |Cx|) \subset P(C)$.

One may represent a circular projection explicitly by a formula. If $C = O$, $\omega = \text{pr}_{P^\perp(C)}(x) / \|\text{pr}_{P^\perp(C)}(x)\|$ and $x \notin P^\perp(O)$ then

$$f_{P(O)}(x) = \frac{\|x\|}{\sqrt{x^2 - (\omega, x)^2}} [x - (\omega, x)\omega]. \quad (2)$$

The circular projection $f_{P(C)}$ keeps the distance of a point x to C . The circular projection of a ball can be a non-convex body, see [2].

The circular projections onto punctured 1-planes (lines) have more simple 'nature' than ones onto higher dimensional planes; circular projection onto any two lines through C are similar in some sense. Here, one can define a **positive circular projection** $|f_C| : \mathbb{E}^n \rightarrow \mathbb{R}_+$ as $|f_C|(x) := |Cx|$.

For the real Euclidean space $\mathbb{E}^n$, the points $|f_C|(x)$ and x belong to the same half-space with respect to hyperplane $P^\perp(C)$. For the complex or the quaternion space $\mathbb{E}^n$ the intersection (with a line) $P(C) \cap S^{an-1}(C, |Cx|)$ ($a = 2, 4$) is a $(a-1)$ -sphere, and $|f_C|(x)$ is the nearest to x point of this sphere.

Remark 4 In order to generalize the notion of circular projections we consider fibration $\pi_0 : \mathbb{E}^n \setminus (0 \times \mathbb{E}^k) \rightarrow \mathbb{E}^{n-k} \setminus \{0\}$, the fiber $S_0(x) = \pi^{-1}(x)$ is a k -dimensional surface, in particular case, such construction has ellipsoidal fibers.

Let $g : [-a, b] \rightarrow \mathbb{R}_+$ (where $0 < a, b \leq \infty$) be a smooth convex function such that $g(0) = 1$, $g'(0) = 0$ and the curvature $k_g \leq 1$. Denote by γ_R a curve $x = Rg(y/R)$ that is either contained in the interior of a real half-space $\mathbb{E}_+^2$ or meets its boundary m transversally.

Suppose that $\mathbb{E}^n = \mathbb{E}^{n-k} \times \mathbb{E}^k$ for some k , and let $\mathbb{E}_+^2$ is orthogonal to both factors (say, x -axis belongs to $\mathbb{E}^{n-k}$ and y -axis belongs to $\mathbb{E}^k$). Now, the group $SO(n)$ contains a subgroup G isomorphic to $SO(n-k) \times SO(k)$ whose action on $\mathbb{E}^n$ splits along the factors $G_1 = SO(n-k)$ and $G_2 = SO(k)$. This gives rise to a G -invariant hypersurface $M_R^{n-1} = G(\gamma_R) \subset \mathbb{E}^n$ called a **hypersurface of revolution with γ_R as profile**. Each M_R^{n-1} intersects $\mathbb{E}^{n-k} \times \{0\}$ by a sphere $G_1(\gamma_R(0)) = S^{n-k-1}(R)$ of radius R , i.e., the G_1 -orbit of a point $\gamma_R(0)$. The action of a subgroup G_2 gives us a fibration $\pi_0 : \mathbb{E}^n \setminus (0 \times \mathbb{E}^k) \rightarrow \mathbb{E}^{n-k} \setminus \{0\}$, the fiber $S_0 = G_2(\gamma_R)$ is the k -dimensional surface. Moreover, M_R^{n-1} is a union of such surfaces.

For example, if $g(y) = \sqrt{1 - x^2}$ then γ_R is a semi-circle and M_R^{n-1} is a sphere of radius R , and $G_1(\gamma_R)$ is a large k -sphere orthogonal to $(n-k)$ -sphere $M_R^{n-1} \cap \mathbb{E}^{n-k}$. Similarly, $g(y) = (b/a)\sqrt{1 - x^2}$ produces the ellipsoids.

Definition 5 Given a punctured $(n-k)$ -plane $P(C)$, consider a Euclidean motion $T : \mathbb{E}^n \setminus P^\perp(C) \rightarrow \mathbb{E}^n \setminus \mathbb{E}^k$. Let $g : [-a, b] \rightarrow \mathbb{R}_+$ be as above. Then $\pi_C := T^{-1} \circ \pi_0 \circ T : \mathbb{E}^n \setminus P^\perp(C) \rightarrow P(C) \setminus C$ is a fibration by k -dimensional surfaces. Denote by $M^k(P(C), x)$ such a fiber through a point $x \notin P^\perp(C)$. For any such point denote by $f_{P(C)}(x)$ the closest to x point of the sphere $S(P(C), g, x) := S(C, R(x)) \cap P(C)$, where $M_{C, R(x)}^{n-1}$ a hypersurface containing x . We call $f_{P(C)} : \mathbb{E}^n \setminus P^\perp(C) \rightarrow P(C) \setminus C$ a **g -non-linear projection** onto $P(C)$. Denote by $S^\perp(P(C), x)$ a k -dimensional 'surface-meridian' in $M_{C, R(x)}^{n-1}$ through x (orthogonal to a sphere $S(P(C), g, x)$).

The definitions (starting from Definition 7) and results in what follows can be generalized for the g -non-linear projections.

2.2 $\mathcal{C}$ -convex bodies and their companions

First, we extend the Definitions 1 and 3.

Definition 6 A ball $B^{k+1}(x, r)$ with the boundary sphere $S^k(x, r)$ in $\mathbb{E}^n$ will be called **supporting** a body $K \subset \mathbb{E}^n$ if $S^k(x, r) \cap K \neq \emptyset$ and $B^{k+1}(x, r)$ is disjoint from $\text{int } K$.

Definition 7 Let $\mathcal{P}_k$ be a non-empty family of circular projections onto punctured $(n-k)$ -planes in $\mathbb{E}^n$. We say that subsets A, B of $\mathbb{E}^n$ are **$\mathcal{P}_k$ - $\mathcal{C}$ -equivalent**, written $A \stackrel{\mathcal{P}_k}{\sim} B$, if and only if $f(A) = f(B)$ for all $f \in \mathcal{P}_k$. For each subset A of $\mathbb{E}^n$ we define $\mathcal{P}_k\langle A \rangle$ to be the union of all subsets B satisfying $B \stackrel{\mathcal{P}_k}{\sim} A$.

The $\mathcal{C}$ -**visual hull** $\mathcal{CP}_k\langle W \rangle$ of a set W in $\mathbb{E}^n$ is the largest set V such that $f(V) \subseteq f(W)$ for all $f \in \mathcal{P}_k$. A body $K \subset \mathbb{E}^n$ is called $\mathcal{P}_k$ - **$\mathcal{C}$ -visual** if $\mathcal{CP}_k\langle K \rangle = K$.

Proposition 4 $\mathcal{CP}_k\langle K \rangle = K$ if and only if for every $x \notin K$ there is $f_{P(C)} \in \mathcal{P}_k$ such that $S^\perp(P(C), x)$ does not intersect K .

Proof almost completely repeats the proof of Proposition 3.

1. Let $x \in \mathcal{CP}_k\langle K \rangle \setminus K$. Thus for all circular projections $f_{P(C)} \in \mathcal{P}_k$ the image of this point, $f_{P(C)}(x)$, belongs to $f_{P(C)}(K)$, but then the corresponding circles $S^\perp(P(C), x)$ will intersect K .

2. Let for any point $x \notin K$ there exists a circular projection $f_{P(C)} \in \mathcal{P}_k$ such that $S^\perp(P(C), x)$ does not intersect K . Thus $f_{P(C)} \notin \mathcal{P}_k$, and we may conclude that any point outside of K does not belong to $\mathcal{CP}_k\langle K \rangle$. $\square$

Consider several useful families $\mathcal{P}_k$.

Example 3 Given a set $M \subset \mathbb{E}^n$, denote by $\mathcal{P}_{k,M}$ the family of circular projections onto punctured $(n-k)$ -planes $P(C)$ in $\mathbb{E}^n$ such that $C \in M$.

If a body K belongs to a ball $B(x, r)$, one may consider $M = S^{n-1}(x, r) -$ a sphere.

The following two examples are used in our study:

1. Given a body $K \subset \mathbb{E}^n$, denote by $\mathcal{P}_{k,K,\varepsilon}$ a family of circular projections onto punctured $(n-k)$ -planes $P(C)$ such that $P^\perp(C) \cap K = \emptyset$, and $\text{dist}(C, K) \leq 1/\varepsilon$. If we take $M = \{C \in \mathbb{E}^n : \text{dist}(C, K) \leq 1/\varepsilon\}$ then we obtain the classes of Definition 8 in what follows.

2. Given a k -plane $L^k \subset \mathbb{E}^n$, denote by $\mathcal{P}_{L^k}$ a family of circular projections onto punctured $(n-k)$ -planes $P(C)$ in $\mathbb{E}^n$ such that $C \in L^k$ and $P(C) \perp L^k$. For $\mathcal{P}_k = \mathcal{P}_{L^k}$ in Definition 8 we obtain 4 classes of bodies, that will be denoted by $\mathcal{KP}_{k,i}(L^k), \mathcal{VP}_{k,i}(L^k), i = 1, 2$.

In order to study the **question** (see Introduction), we will extend the Definition 2.

Definition 8 A body $K \subset \mathbb{E}^n$ is called $\mathcal{C}$ -**convex** of a class $\mathcal{K}_i^{k,\varepsilon}$ (for some $\varepsilon > 0$ and $0 \leq k \leq n-1$) if

$\mathcal{K}_1^{k,\varepsilon}$: for any point $x \in \partial K$ there is a $f_{P(C)} \in \mathcal{P}_{k,K,\varepsilon}$ such that $x \in P(C)$, $|Cx| = 1/\varepsilon$ and $f_{P(C)}(x) \in \partial(f_{P(C)}(K))$,

$\mathcal{K}_2^{k,\varepsilon}$: for any point $x \notin K$, $\text{dist}(x, K) \leq 1/\varepsilon$, there is a $f_{P(C)} \in \mathcal{P}_{k,K,\varepsilon}$ such that $x \in P(C)$, $|Cx| = 1/\varepsilon$ and $f_{P(C)}(x)$ is disjoint from $f_{P(C)}(K)$.

A body $K \subset \mathbb{E}^n$ is called $\mathcal{C}$ -**visible** of a class $\mathcal{V}_i^{k,\varepsilon}$ if

$\mathcal{V}_1^{k,\varepsilon}$: for any ball $\tilde{B}^{k-m+1}(C, 1/\varepsilon)$ supporting K there is a $f_{P(C)} \in \mathcal{P}_{k,K,\varepsilon}$ such that $\partial \tilde{B} \perp S(P(C), 1/\varepsilon)$ and $f_{P(C)}(\tilde{B})$ is supporting $f_{P(C)}(K)$,

$\mathcal{V}_2^{k,\varepsilon}$: for any sphere $\tilde{S}^{k-m}(C, 1/\varepsilon)$ disjoint from K and $\text{dist}(C, K) \leq 1/\varepsilon$, there is a $f_{P(C)} \in \mathcal{P}_{k,K,\varepsilon}$ such that $\tilde{S} \perp S(P(C), 1/\varepsilon)$ and $f_{P(C)}(\tilde{S})$ is disjoint from $f_{P(C)}(K)$.

Here, as above, $0 < m \leq k < n$, and the 0-dimensional sphere is a point.

Similarly one may define the classes $\mathcal{KP}_{k,i}$ ($i = 1, 2$) of $\mathcal{C}$ -**convex** and $\mathcal{C}$ -**visible** bodies in $\mathbb{E}^n$ relative to **any** family $\mathcal{P}_k$ (in particular, for $\mathcal{P}_{L^k}$ and $\mathcal{P}_{k,M}$) of circular projections onto punctured $(n-k)$ -planes $P(C)$.

One may verify (applying just the set theory arguments) that if a body K is the intersection of connected bodies of a class $\mathcal{K}_2^{k,\varepsilon}$ then K also belongs to $\mathcal{K}_2^{k,\varepsilon}$. This intersection can be disconnected.

Proposition 5 *The conditions $K \in \mathcal{K}_i^{k,\varepsilon}$ ($i = 1, 2$) are equivalent to the following ones:*

- ($i = 1$) *any point $x \in \partial K$ belongs to a ball $B^{k+1}(C, 1/\varepsilon)$ supporting K ,*
- ($i = 2$) *any point $x \notin K$, $\text{dist}(x, K) \leq 1/\varepsilon$, belongs to the boundary sphere of a ball $B^{k+1}(C, 1/\varepsilon)$ disjoint from K .*

The condition $K \in \mathcal{V}_i^{k,\varepsilon}$ ($i = 1, 2$) is equivalent to the following ones:

- ($i = 1$) *any ball $B^{k-m+1}(C, 1/\varepsilon)$ supporting K belongs to a ball $B^{k+1}(C, 1/\varepsilon)$ supporting K as well,*
- ($i = 2$) *any sphere $S^{k-m}(C, 1/\varepsilon)$ disjoint from K and $\text{dist}(C, K) \leq 1/\varepsilon$ belongs to the boundary sphere of a ball $B^{k+1}(C, 1/\varepsilon)$ disjoint from K as well.*

The proof of Proposition 5 obviously follows from the definitions.

Remark 5 We have introduced in [2] a " $k = n - 1$ version" of the $\mathcal{C}$ -convex objects. A body $K \subset M$ was called ε -**convex** of a class $\mathcal{K}_i^\varepsilon$ (for some $\varepsilon > 0$) if

$\mathcal{K}_1^\varepsilon$: *any point $x \in \partial K$ belongs to a supporting ball (of K) of radius $1/\varepsilon$,*

$\mathcal{K}_2^\varepsilon$: *any point $x \notin K$ belongs to a ball B of radius $1/\varepsilon$ such that $\text{int}K \cap B = \emptyset$.*

Here only the class $\mathcal{K}_1^\varepsilon$ coincides with $\mathcal{K}_1^{n-1,\varepsilon}$. Simple examples show that corresponding classes $\mathcal{K}_2^{n-1,\varepsilon}$ and $\mathcal{K}_2^\varepsilon$ are different.

Theorem 2 *The strong inclusions hold for all $\varepsilon > 0$, $k < n$, $i = 1, 2$:*

$$\mathcal{K}_2^{k,\varepsilon} \subset \mathcal{K}_1^{k,\varepsilon}, \quad \mathcal{V}_2^{k,\varepsilon} \subset \mathcal{V}_1^{k,\varepsilon}, \quad \text{and} \quad \mathcal{V}_i^{k,\varepsilon} \subset \mathcal{K}_i^{k,\varepsilon}.$$

Proof is similar to the proof of Theorem 1 in [2].

$\mathcal{K}_2^{k,\varepsilon} \subset \mathcal{K}_1^{k,\varepsilon}$. Let $x_* \in \partial K$, $K \in \mathcal{K}_2^{k,\varepsilon}$. Consider a sequence of points $x_i \notin K$ converging to x_* . By condition, for any i there is a sphere $S^k(C_i, 1/\varepsilon)$, containing x_i and disjoint from K , moreover, $\text{dist}(C_i, K) \leq 1/\varepsilon$. The bounded sequence of points $\{C_i\}$ has a limit point C_* , moreover, $\text{dist}(C_*, K) \leq 1/\varepsilon$ and

$$1/\varepsilon \geq \lim_{i \rightarrow \infty} \rho(x_i, C_i) = \rho(x_*, C_*).$$

Hence, $x_* \in S^k(C_*, 1/\varepsilon)$ and a sphere $S^k(C_*, 1/\varepsilon)$ is disjoint from $\text{int}K$ and determines a circular projection from a class $\mathcal{K}_1^{k,\varepsilon}$.

Other inclusions can be proved analogously. $\square$

Example 4 One can produce examples of bodies $K \in \mathcal{K}_1^{k,\varepsilon} \setminus \mathcal{K}_2^{k,\varepsilon}$ applying small deformations to corresponding examples of $K \in \mathcal{K}_1^k \setminus \mathcal{K}_2^k$, see Fig. 1, where a circular domain about the coordinate center is replaced by a spherical shell of radius $1/\varepsilon$, see Fig. 2.




Figure 2: $K \in \mathcal{K}_1^{k,\varepsilon} \setminus \mathcal{K}_2^{k,\varepsilon}$ ($k = 1$).

The following proposition was proved in [2] for $k = n - 1$.

Proposition 6 *If a connected body $K \in \mathcal{K}_1^{k,\varepsilon}$ has $\text{diam } K < 1/\varepsilon$ then the boundary ∂K is connected as well.*

Proof. Let K be connected and its boundary ∂K is not, then one of components of ∂K , denote it by $\partial_1 K$, separates K from "infinity" (or from $\mathbb{E}^n \setminus \text{conv } K_\varepsilon$). Now, $\partial_1 K$ is a boundary of some body L such that $\text{diam } L = \text{diam } (\partial K) = \text{diam } K$. Let $\partial_2 K \neq \partial_1 K$ be another component of ∂K and let $x_1 \in \partial_2 K$. If $K \in \mathcal{K}_1^{k,\varepsilon}$ then x_1 should be contained in L and in some supporting ball $B^{k+1}(C, 1/\varepsilon)$, moreover, $\text{dist}(C, K) \leq 1/\varepsilon$. So, $B^{k+1}(C, 1/\varepsilon) \subset L$ and $\text{diam } L \geq 1/\varepsilon$, a contradiction. $\square$

Next we consider a special case of Definition 7.

Definition 9 The ε -visual hull $\mathcal{CP}_{k,\varepsilon}\langle V \rangle$ of a set V in $\mathbb{E}^n$ (defined by the family $\mathcal{P}_{k,V,\varepsilon}$ of circular projections) is the largest set K such that $f(K) \subseteq f(V)$ for all $f \in \mathcal{P}_{k,V,\varepsilon}$. A body $K \subset \mathbb{E}^n$ is called ε -visual (relative to $\mathcal{P}_{k,K,\varepsilon}$) if $\mathcal{CP}_{k,\varepsilon}\langle K \rangle = K$.

Proposition 7 *If $\text{diam } K < 1/\varepsilon$, then $K \in \mathcal{K}_2^{k,\varepsilon}$ if and only if $\mathcal{CP}_{k,\varepsilon}\langle K \rangle = K$.*

Proof. From definition of the class $\mathcal{K}_2^{k,\varepsilon}$ it follows that for any point $x \notin K$ there is a circular projection $f_{P(C)} \in \mathcal{P}_{k,K,\varepsilon}$ such that $|Cx| = 1/\varepsilon$ and $f_{P(C)}(x)$ is disjoint from $f_{P(C)}(K)$. Hence, any point outside of K does not belong to its ε -visual hull $\mathcal{CP}_{k,\varepsilon}\langle K \rangle$. $\square$

Remark 6 For the "large" bodies ($\text{diam } K \geq 1/\varepsilon$) this proposition is wrong:

Let $K = B(0, 3/\varepsilon) \setminus B(0, 2/\varepsilon)$. For any point $C \in B(0, 2/\varepsilon)$ and for any plane $P(C)$ of any dimension the orthogonal complement $P^\perp(C)$ intersects this body K which obviously belongs to $\mathcal{K}_2^{k,\varepsilon}$, for any k , $1 < k < n$ (for "small" bodies, i.e., $\text{diam}(K) < 1/\varepsilon$, this claim fails). Hence, for determination of the ε -visual hull $\mathcal{CP}_{k,\varepsilon}(K)$ one should take the centers C outside of the "large" ball $B(0, 3/\varepsilon)$, i.e., we have here $\mathcal{CP}_{k,\varepsilon}(K) = B(0, 3/\varepsilon)$.

2.3 Reconstruction of $\mathcal{C}$ -convex bodies by their circular projections

Theorem 3 *If the circular projections of $V_1, V_2 \in \mathcal{K}_2^{n-1, \varepsilon}$ onto any punctured line $P^1(C)$ coincide, and $\text{diam} V_1, \text{diam} V_2 < 1/(2\varepsilon)$, then $V_1 = V_2$. Here C is a center of any ball of radius $1/\varepsilon$ supporting V_1 or V_2 .*

Proof. 1. It follows from the conditions that if $B(C, 1/\varepsilon)$ is supporting V_1 , then it is supporting to V_2 and vice versa. In the same way one can verify that $\text{dist}(V_1, V_2) < 1/(2\varepsilon)$.

2. Hence $\text{conv}(V_1) = \text{conv}(V_2)$. We denote it by K , its diameter is less $1/(2\varepsilon)$ as well and any center of supporting ball with radius $1/\varepsilon$ can not be contained in K .

3. Now, if $y_2 \in V_2 \subset K$ and $y_2 \notin V_1$, then we can assume that $y_2 \in \text{int} V_2$; $y_2 \in B(C(y_2), 1/\varepsilon) \equiv B_2$, B_2 is disjoint from $\text{int} V_1$ by definition of the class $\mathcal{K}_2^{n-1, \varepsilon}$. This ball B_2 can not be supporting V_1 , otherwise it should be supporting to V_2 . So, B_2 is disjoint from V_1 .

4. Let $E(y_2)$ be the equatorial hyperplane of B_2 orthogonal to the line $[C(y_2), y_2]$. This hyperplane divides the space $\mathbb{E}^n$. Since the bodies V_1, V_2 are "small", one of these half-spaces $\mathbb{E}_+^n$, contains V_1 and V_2 , another one $\mathbb{E}_-^n$ does not contain them.

5. We shall move B_2 in the direction $\overrightarrow{y_2, C(y_2)}$ – outside of K and the result will be denoted by $B_2(t)$. During this displacement the balls $B_2(t)$ will be contained in the union $B_2 \cup \mathbb{E}_-^n$, so they will never intersect V_1 . At some moment t_1 the ball $B_2(t_1)$ will be supporting V_2 . But when B_2 moves outside of K it can not touch V_1 . This contradiction follows from assumption 3.

Hence $V_1 = V_2$. $\square$

Remark 7 For "large" bodies V_1, V_2 , $\text{diam} V_1, \text{diam} V_2 > 1/\varepsilon$ and for positive circular projection Theorem 3 is wrong:

Let $V_1 \subset \mathbb{E}^2$ be a ring $1 + 2/\varepsilon \leq |x| \leq 3 + 2/\varepsilon$ and $V_2 = V_1 \cup B(0, 1)$, where $B(0, 1)$ is a unit ball centered in the origin. All the balls $B(Z, 1/\varepsilon)$ centered in Z , $|Z| = 1 + 1/\varepsilon$ are supporting both for V_1 and V_2 (and for $B(0, 1)$). Clearly, for all such Z we have $|f_Z|(V_1) = |f_Z|(V_2)$.

Theorem 3 also fails for a wider class $\mathcal{K}_1^{n-1, \varepsilon}$ that can be seen from examples similar to above one.

Proposition 8 *If a body $V_1 \subset \mathbb{E}^n \setminus L^k$ belongs to a class $\mathcal{KP}_{k,2}(L^k)$ ($1 \leq k \leq n-1$), and V_2 is obtained from V_1 by some rotation of $\mathbb{E}^n$ about L^k (i.e., by some transformation in $SO(n-k)$), then for any $C \in L^k$ the circular projections $f_{P(C)}$ of the bodies V_1 and V_2 onto the plane $P^{n-k}(C)$ are congruent with respect to some rotation of this plane (or are $SO(n-k)$ -equivalent).*

Proof. The proof immediately follows from the fact that for any $C \in L^k$ the sphere $S^{n-1}(C, r)$ and decomposition of the space $\mathbb{E}^n = L^k \oplus P^{n-k}(C)$ are invariant with respect to the action of $SO(n-k)$. $\square$

We will say that a body $K \subset \mathbb{E}^n$ **surrounds** a plane P^{n-2} if any half-plane H^{n-1} bounded by P^{n-2} intersects K .

Here is an example of an answer to the **question** from the Introduction.

Theorem 4 *If the bodies $V_1, V_2 \subset \mathbb{E}^n \setminus L^{n-2}$ belong to a class $\mathcal{KP}_{n-2,2}(L^{n-2})$ and their circular projections onto all planes $P^2(C)$ ($C \in L^{n-2}$) orthogonal to L^{n-2} are $SO(2)$ -equivalent, then V_1 can be obtained from V_2 by some rotation about L^{n-2} .*

Proof of Theorem 4. For visuality, we divide the proof into two steps: 3-dimensional and multidimensional. Let $n - 2 = 1$.

1. Fix some $C_0 \in L^1$ and rotate V_1 about the axis L^1 so that the image V'_1 and the body V_2 have equal circular projection onto $P^2(C_0)$, i.e., $f_{C_0}(V'_1) = f_{C_0}(V_2)$. These projections are contained in some angle $A_0C_0B_0$ in the plane $P(C_0)$ and touch the edges of this angle. Hence, the bodies V'_1 and V_2 belong to the dihedral angle $A_0C_0B_0$ with the edge L^1 and touch the faces of this angle.

2. If for some $C_1 \in L^1$ the circular projections $f_{C_1}(V'_1)$ and $f_{C_1}(V_2)$ do not coincide and are equivalent with respect to rotation in the plane $P(C_1)$ by some angle $\phi(C_1)$, $0 < \phi(C_1) < 2\pi$, then $f_{C_1}(V'_1)$ is contained in some plane angle $A_1C_1B_1$ and touches its edges, and the circular projection $f_{C_1}(V_2)$ is contained in some equivalent angle $A'_1C_1B'_1$ and touches its edges. Hence, the bodies V'_1 and V_2 are contained in dihedral angles $A_1C_1B_1$, $A'_1C_1B'_1$, respectively; these angles have common edge L^1 and their faces touch these bodies V'_1 and V_2 . This contradicts to the existence of the dihedral angle $A_0C_0B_0$ constructed above.

Now let $n - 2 > 1$.

3. Fix as above any $C_0 \in L^{n-2}$ and rotate the body V_1 about the axis L^{n-2} so that the image V'_1 and the body V_2 have equal circular projection $f_{C_0}(V'_1) = f_{C_0}(V_2)$ in $P^2(C_0)$.

4. If for some $C_1 \in L^{n-2}$ the circular projections $f_{C_1}(V'_1)$ and $f_{C_1}(V_2)$ do not coincide, but are equivalent with respect to some rotation $\varphi(C_1) \in SO(2)$ of the plane $P^2(C_1)$, then since the bodies V_1, V_2 do not surround L^{n-2} , their circular projections $V'_1(P^2(C_1))$ and $V_2(P^2(C_1))$ as in 3-dimensional case, belong to equal plane angle $A_1C_1B_1$ and $A'_1C_1B'_1$ with a common $(n - 2)$ -dimensional edge L^{n-2} and touch their edges. As above, the bodies V'_1, V_2 are contained in corresponding dihedral angles $A_1C_1B_1$ and $A'_1C_1B'_1$ with common $(n - 2)$ -dimensional "edge" L^{n-2} and touch their faces.

These angles are congruent with respect to the rotation by the angle $\varphi(C_1)$ in the plane $P^2(C_1)$. On the other hand the intersections of the faces with the plane $P^{n-k}(C_0)$ generate in that plane the dihedral angles with common vertex C_0 . These dihedral angles contain the circular projections $f_{C_0}(V'_1) = f_{C_0}(V_2)$ and they are congruent with respect to the same nontrivial rotation by the angle $\varphi(C_1)$, a contradiction to non-triviality of this angle. $\square$

Remark 8 (a) The case of $SO(2)$ -equivalent orthogonal projections on all 2-planes, where under additional assumption that the projections of bodies have no $SO(2)$ -symmetries, it was proven that the bodies V_1, V_2 are equal with respect

either translation or central symmetry, was studied in [3]. In contrast to this study, the conditions of Theorem 4 do not contain asymmetry assumptions, but

- only orthogonal to L^{n-2} planes of circular projection are considered,
- the centers of these circular projections belong to L^{n-2} .

(b) In the same way Theorem 4 can be reformulated for the cases of complex and quaternion Euclidean spaces where one should consider $U(n-k)$, $SU(n-k)$ and $Sp(n-k)$ -transformations. In [8] corresponding results were obtained for the case of orthogonal projections onto $(n-k)$ -planes of complex Euclidean space for arbitrary k and under assumption that these projections have no $U(n-k)$ or $SU(n-k)$ symmetries.

References

- [1] Yu. G. Reshetnyak, On one generalization of convex surfaces. Math. Sbornik, 1956, 40 (3), 381–398 (Russian).
- [2] V. P. Golubyatnikov, V. Y. Rovenski, Notes on ε -convex bodies, submitted to Israel J. of Math. (2008), 1–18.
- [3] V. P. Golubyatnikov, Uniqueness Questions in Reconstruction of Multi-dimensional Objects from Tomography-Type Projection Data. VSP BV, The Netherlands, 2000.
- [4] G. H. Meisters, S. M. Ulam, On visual hull of sets. Proc. Nat. Acad. Sci. USA, 1967, 57, 1172–1174.
- [5] D. G. Larman, P. Mani, On visual hulls. Pacific J. Math., 1970, 32(1), 157–171.
- [6] S. Shefel', About two classes of k -dimensional submanifolds in n -dimensional Euclidean space. Siberian Math. J., 1969, 10(2): 459–467.
- [7] V. P. Golubyatnikov, Stability problems in certain inverse problems of reconstruction of convex compacta from their projections (English). Siberian Math. J. 33(3), (1992), 409–415.
- [8] V. P. Golubyatnikov, On the unique determination of compact convex sets from their projections. The complex case. Siberian Math. J. 40(4), (1999), 678–681.
- [9] V. Yu. Rovenski, On (k, ε) -saddle submanifolds of Riemannian manifolds. Geometriae Dedicata, 121 (2006), 187–203.
- [10] M. Gromov, Spaces and questions. Geom. Funct. Anal. (GAFA), Special Volume, Part I (2000), 118–161.
- [11] R. Gardner, Geometric Tomography, Cambridge University Press, 2006.
- [12] A. A. Borisenko, On the extrinsic geometric properties of parabolic surfaces and topological properties of saddle surfaces in symmetric spaces of rank one. Math. USSR Sb., 42: 297–310, 1982.
- [13] S. Lazebnik, Y. Furukawa and J. Ponce, Projective visual hulls. Int. J. of Computer Vision, 2007, 74(2), 137–165.